



DO YOU HAVE PAD?

Common Symptoms of Peripheral Arterial Disease (PAD):

- Intermittent claudication of the legs
- Leg and muscle weakness
- Numbness or tingling sensation
- Weak pulse in legs
- Cold extremities
- Non-healing wounds



1 in 3 Americans over the age of 50 with diabetes are likely to have PAD.



PAD affects a shocking **12-20%** of Americans age 65 or older.



Untreated, PAD can lead to **CLTI (Critical Limb Threatening Ischemia)** and **amputation**.



People with PAD have a **4-5x GREATER RISK** of a heart attack or a stroke.

EARLY IDENTIFICATION IS CRITICAL

Early identification and management of patients with **PAD** can **improve quality of life, save limbs & reduce cardiovascular events.**

**AT RISK? TALK TO
YOUR DOCTOR.**



DO YOU HAVE PAD?

Important information about Peripheral Arterial Disease

SIGNS & SYMPTOMS OF PAD

- Intermittent claudication of the legs
- Leg and muscle weakness
- Weak pulse in legs
- Cold extremities
- Non-healing wounds
- Numbness or tingling sensation



1 in 3 Americans over 50 with diabetes are likely to have PAD.

PAD affects **12-20%** of Americans age 65 or older.

Left untreated, PAD can lead to **CLTI (Critical Limb Threatening Ischemia)** and **amputation**.

EARLY IDENTIFICATION IS CRITICAL

ARE YOU AT RISK? TALK TO YOUR DOCTOR



1 in 3 Americans over age 50 with diabetes are likely to have PAD.

Early assessment and management of patients with PAD can improve quality of life, save limbs & reduce cardiovascular events.